arXiv:1602.02534v1 [astro-ph.GA] 8 Feb 2016

High-Resolution Imaging of Water Maser Emission in the active galaxies NGC 6240 and M51

Yoshiaki Hagiwara[1] and Philip G. Edwards[2]
[1] *Natural Science Laboratory, Toyo University, 5-28-20, Hakusan, Bunkyo-ku, Tokyo 112-8606, Japan; yhagiwara@toyo.jp*
[2] *CSIRO Astronomy and Space Science, P.O. Box 76, Epping NSW 1710, Australia; Philip.Edwards@csiro.au*
Received 2015 September 23; accepted 2015 November 10; published 2015 December 17

ABSTRACT

We present the results of observations of 22 GHz H_2O maser emission in NGC 6240 and M51 made with the Karl G. Jansky Very Large Array. Two major H_2O maser features and several minor features are detected toward the southern nucleus of NGC 6240. These features are redshifted by about 300 km s^{-1} from the galaxy's systemic velocity and remain unresolved at the synthesized beam size. A combination of our two-epoch observations and published data reveals an apparent correlation between the strength of the maser and the 22 GHz radio continuum emission, implying that the maser excitation relates to the activity of an active galactic nucleus in the southern nucleus rather than star-forming activity. The star-forming galaxy M51 hosts H_2O maser emission in the center of the galaxy; however, the origin of the maser has been an open question. We report the first detection of 22 GHz nuclear radio continuum emission in M51. The continuum emission is co-located with the maser position, which indicates that the maser arises from active galactic nucleus-activity and not from star-forming activity in the galaxy.

Subject headings: galaxies: active - galaxies: individual (NGC6240, M51) - galaxies: ISM - galaxies: nuclei - masers

1. Introduction

Observations of 22 GHz H_2O masers have become established as a powerful tool for investigating the inner few parsecs of active galactic nuclei (AGNs), and have enabled precise distance measurements to local group galaxies and nearby AGNs (e.g., Herrnstein et al. 1999; Greenhill et al. 2009). Observations of extragalactic masers are of interest for determining the origin of the maser emission (e.g., whether it is driven by AGNs or starburst activity), and the suitability of the galaxy for direct distance measurement. Extragalactic H_2O masers have also been detected in the sub-millimeter bands, with the 183 GHz, and possibly 439 GHz, transitions detected in NGC 3079 (Humphreys et al. 2005), and the 321 GHz transition detected in the Circinus galaxy (Hagiwara et al. 2013), providing additional information on the physical environments hosting these masers. In this paper we describe the results of observations with the Karl G. Jansky Very Large Array (VLA) of lower-luminosity 22 GHz H_2O maser emission in the galaxies NGC 6240 (J1652+0024) and M51 (J1329+4713).

NGC 6240 is an interacting/merging galaxy hosting two nuclei that are separated by $\sim$1.″5 (Beswick et al. 2001). At 22 GHz, an H_2O maser with low isotropic luminosity ($L_{H_2O} \lesssim 10\ L_\odot$) has been reported in this galaxy (Henkel et al. 1984; Hagiwara et al. 2002; Nakai et al. 2002). NGC 6240 has a large far-infrared luminosity $\approx L_{FIR} = 10^{11-12}\ L_\odot$, which cannot be accounted for by starburst activity alone, but which can be more naturally explained as a buried AGN heating surrounding dusty com-

ponents (e.g., Depoy, Becklin, & Wynn-Williams 1986; Sanders et al. 1988; Risaliti et al. 2006). Neutral Fe Kα lines at 6.4 keV due to reflection from optically thick material are detected from both nuclei, which is clear evidence for both being AGNs (Komossa et al. 2003). Very Long Baseline Interferometry (VLBI) observations (Gallimore & Beswick 2004; Hagiwara et al. 2011) detected compact radio sources with brightness temperatures of 7×10^6 K for the northern nucleus and 1.8×10^7 K for the southern nucleus (e.g., Colbert et al. 1994), confirming the presence of AGNs at centimeter radio wavelengths. Using the VLA, we found that the maser emission was coincident with the southern nucleus to within the $0.''1$ resolution of the observation (Hagiwara et al. 2003). The origin of the maser has been an open question for some time (Hagiwara 2010).

The star-forming galaxy M51 (NGC 5194) hosts a Seyfert/LINER nucleus (Heckman 1980) and has H_2O maser emission (Ho et al. 1987) that consists of Doppler-shifted features on either side of the galaxy's systemic velocity (Hagiwara et al. 2001). The redshifted features are located toward the nuclear radio source, while the location of the blue-shifted counterpart is displaced from the radio source toward the southeast. A small velocity gradient closely aligned with the radio jet is detected from the redshifted features (Hagiwara 2007). The redshifted maser most likely amplifies the background radio continuum jet, while the blue-shifted counterpart marks off-nuclear star formation in the galaxy.

The upgraded VLA capabilities, with bandwidths of 2 GHz and improved spectral resolution (Perley et al. 2011), have allowed us to address the nature of the low-luminosity H_2O masers in NGC 6240 and M51 and to map the distribution of maser spots on the 0.1 arcsec scale.

We adopt cosmological parameters of $H_0 = 73$ km s^{-1} Mpc^{-1}, $\Omega_\Lambda = 0.73$, and $\Omega_M = 0.27$. The luminosity distance to NGC 6240 is therefore 103 Mpc (z=0.02448) and 10.6 Mpc (z=0.002) to M51. Thus, 1 arcsec corresponds to a linear scale of 475 pc for NGC6240 and 51 pc for M51. We adopt the optical velocity definition and velocities are calculated with respect to the local standard of rest (LSR).

2. OBSERVATIONS AND DATA REDUCTION

We observed NGC 6240 with the National Radio Astronomy Observatory (NRAO)[1] VLA in the B configuration on 2012 July 21 (epoch 1) and 2012 August 28 (epoch 2) and observed M 51 in the A configuration on 2012 November 17. Observations were made of 22 GHz H_2O maser emission in the $6_{16}-5_{23}$ transition, employing 16 intermediate frequencies (IFs) of 16 MHz bandwidth with dual polarization, resulting in a total bandwidth of 256 MHz for each polarization. The total velocity coverage is $\sim$3450 km s^{-1}, however, the velocity coverages were not continuous in both observations: there is a $\sim$430 MHz frequency gap between IFs 1–8 and IFs 9–16, where IF4 is centered on the systemic velocity of each galaxy. The total observing time for NGC 6240 at each epoch was $\sim$2.5 hr and that of M51 was $\sim$3 hr. In the NGC 6240 observations, the phase-referencing observations were executed in a sequence of 3 minute scans with 2 minutes on NGC 6240 and 40 s on J1658+0741. For the M51 observation, the phase-referencing was performed in a sequence of 3.75 minute scans with 2 minutes on M51 and 40 s on J1419+5423. Flux density and bandpass calibration were performed using observations of 3C 286. Standard VLA data calibration and editing were made using the NRAO Astronomical Image Processing Software (AIPS). Uncertainties of flux density were estimated in the same method as explained in Hagiwara (2010).

After the phase and flux calibrations and data editing, the continuum emission was subtracted from the spectral-line visibilities using line-free channels prior to the imaging and CLEAN deconvolution of the maser emission. Imaging and CLEANing of the continuum were done separately.

The synthesized beam sizes produced from naturally weighted spectral-line images were 0.38×0.34 arcsec^2 (PA=18°) for NGC 6240 and 0.10×0.09 arcsec^2 (PA=-3°) for M51. The spectral resolution was 125 kHz (1.7 km s^{-1}) per spectral channel for all observations. The rms noise levels of the naturally weighted images at 125 kHz resolution were $\sim$1.5$-$2.0 mJy

[1]The NRAO is operated by Associated Universities, Inc., under a cooperative agreement with the National Science Foundation.

beam^{-1}(epoch 1), $\sim$0.8$-$1.3 mJy beam^{-1}(epoch 2) for NGC 6240 and $\sim$1.0$-$1.2 mJy beam^{-1} for M51. The rms noises of the continuum images were about 0.14 mJy beam^{-1} for NGC 6240, and 0.036 mJy beam^{-1} for M51, respectively.

3. RESULTS

3.1. NGC 6240

Figure 1 shows two-epoch spectra of the H_2O maser emission from the southern nucleus in the velocity range spanning $V_{\rm LSR}$=7450$-$7690 km s^{-1}. The total velocity coverages in our observations are $V_{\rm LSR}\approx$ $-$730$-$990 km s^{-1} and $\approx$ 6590 – 8320 km s^{-1}. The maser line peaking at $V_{\rm LSR}$=7609.3 km s^{-1} is clearly detected at both epochs, though it is somewhat weaker at the second epoch. This location of this feature was determined with the VLA in 2012 to be coincident with the southern nucleus. A marginally significant feature at $V_{\rm LSR}$=7563.2 km s^{-1} in the first epoch matches features seen by single-dish measurements in 2000$-$2001, and 2005 (see Table 2), but there is no evidence of this feature in the second epoch.

Figure 2 shows the spectra of tentative features peaking at $V_{\rm LSR}$=7158km s^{-1} , 7372 km s^{-1} , and 7396 km s^{-1} and in the southern nucleus. The maser emission also exhibits a broad profile that ranges from $V_{\rm LSR}\approx$ 7360 to 7400 km s^{-1}. These features are detected, tentatively, for the first time in the galaxy. No other H_2O maser is detected at either epoch above the 3σ level of $\sim$3 mJy beam^{-1} in the spectra from the northern nucleus, or at any other point within a field of view 7.7$''$ $\times$ 7.7$''$, centered on the southern nucleus.

Figure 3 shows the 22 GHz continuum image from epoch 1 data, obtained by integrating over the full frequency range after the removal of the maser line emission. The image shows two major radio sources, the southern and northern nucleus, both coinciding with the radio sources at lower frequencies with similar angular resolution (e.g., Carral et al. 1990; Colbert et al. 1994; Beswick et al. 2001). Extended components seen in lower frequency VLA images in the literature (e.g., Colbert et al. 1994; Baan, Hagiwara & Hofner 2007) were resolved out in our continuum image. The flux densities of the 22 GHz continuum of the two nuclei and H_2O maser are summarized in Table 1.

All of the masers detected in our observations remain unresolved at the angular resolution of $\sim$ 0.$''$2, or 100 pc. The position of the 7609 km s^{-1} feature is α(J2000): 16$^{\rm h}$52$^{\rm m}$58$^{\rm s}$.886, δ(J2000): +02°24$'$03.$''$260, from which all other features reside within $\approx$ 0.1$''$, about a half of the synthesized beam size of our observation. Positional accuracy, without consideration of systematic errors, is $\sim$ $\theta_{\rm beam}/2\times$S/N $\sim$ 0$''$.02, where $\theta_{\rm beam}$ is the synthesized beam of our observations and S/N is the signal-to-noise ratio of the maser spectra. The position uncertainties considering the position error of the calibrator source (Reid and Honma 2014) are estimated to be $\Delta\alpha \sim 0.^s02$ and $\Delta\delta \sim 0.036''$.

The positions of these masers coincide with that of the southern nucleus, which is consistent with results of the previous VLA observations (Hagiwara 2010). The estimation of the relative positional error between the maser and the southern continuum peak ($\sim$3 pc or less) is considered in Hagiwara et al. (2003) and Hagiwara (2010). A cumulative list of detected H_2O maser features in the galaxy, which is revised from the table 2 in Hagiwara (2010), is presented in table 2.

Figure 4 shows the time variability of the maser and nuclear continuum flux densities, taking three-epoch VLA data obtained from 2002–2009 (see Figure 3 in Hagiwara (2010). There is a similar decrease, in absolute terms, in the flux densities of the southern nucleus continuum emission and the water maser emission over one decade. With two epochs in 2012 and three earlier epochs, the result is strongly suggestive, but further monitoring is required before a conclusive claim for correlation can be made.

3.2. M51

Figure 5 shows the spectrum of the maser at $V_{\rm LSR}$=453.4$-$668.2 km s^{-1}. No maser has been detected outside this range, with the full range searching covering $V_{\rm LSR}\approx$ $-$400 – 1300 km s^{-1}. The detected maser features are redshifted with respect to the galaxy's systemic velocity of $V_{\rm LSR}$=472$\pm$3km s^{-1} (Scoville & Young 1983). All of the detected maser emission remains unresolved at the synthesized beam size of 0.1 arcsec. The positions of the features in the spectrum are all confined within $\approx$ 0.01$''$ from

the position of a feature at $V_{\rm LSR}$= 565.1 km s^{-1} : α (J2000) = $13^{\rm h}29^{\rm m}$ $52.708^{\rm s}$, δ (J2000) = $+47°11'42.810''$, which is consistent with those of earlier measurements in the VLA A configuration (Hagiwara et al. 2001; Hagiwara 2007).

A naturally weighted 22 GHz continuum image of M51 is shown in Figure 5. The continuum emission has been detected with an S/N of ∼6. This is the first reported detection of the 22 GHz radio continuum in M51. The image presents a nuclear source located at α(J2000) = $13^{\rm h}29^{\rm m}52\overset{\rm s}{.}708$, δ(J2000) = $+47°11'42.812''$, where the relative errors between the continuum source and the maser are ∼ 0".01–0".025). Thus, the positions of the unresolved maser sources and continuum emission are located in the same position in the galaxy within errors. The continuum image shows a jet component that is nearly northeast at PA = 158 ± 19°, the axis of which is consistent with that on the lower frequency VLA images (Bradley et al. 2004; Dumas et al. 2011). Table 3 provides a summary of radio flux densities and the spectral index of the radio source. In the VLA full-track observation performed in 2003 July, a weak velocity gradient from $V_{\rm LSR}$=554 to 563 km s^{-1} was identified at PA=155°, nearly along the nuclear radio jet (Hagiwara 2007); however; in our new observation no distinct velocity gradient is apparent. This may be the result of source evolution, though we note that the 2012 epoch had an observing duration of 3 hr as opposed to 8 hr in the 2003 observation, and somewhat poorer (u, v) coverage. The spectral sensitivity does not differ significantly from the previous run conducted in 2003 (Hagiwara 2007).

4. DISCUSSION

4.1. NGC 6240

From two-epoch observations in 2012 and three earlier epochs from 2002 to 2009, we find that the strength of the maser and continuum emission from the AGNs in the southern nucleus vary in a correlated fashion. This strongly suggests that the maser in the galaxy amplifies the background continuum emission, which is exhibiting flux variability that is consistent with that typically observed from the central engine in AGNs. These observations constitute evidence for the maser in NGC 6240 being a nuclear maser similar to the cases of other H_2O megamasers explored to date.

Further observations at VLBI angular resolutions would be required to determine whether the maser is a "disk maser," tracing an inclined disk located at radii ∼0.1–1 pc from the central engine (e.g., Greenhill et al. 2009). It is also interesting to note the tentative detections of some new weaker features in the second epoch, whereas the prominent features at $V_{\rm LSR}$= 7563.2 and 7609.3 km s^{-1} have weakened. However, the emergence of the tentative features at epoch 2 is in contrast with the fact that the 7609 km s^{-1} maser and the 22 GHz continuum emission weakened in epoch 2.

It is possible that these new features are not amplifying the background continuum but are associated with a starburst activity in the center of the galaxy (Genzel et al. 1998) due to the low-luminosity of these features, which is consistent with star-forming activity (Hagiwara 2007). Further monitoring of the velocity ranges in which these new tentative features are seen is required to confirm their existence and trace their variability against the nuclear continuum.

All the maser features reported previously in the galaxy were redshifted by ∼ 200−300 km s^{-1} with respect to both the systemic velocity of the galaxy ($V_{\rm LSR}$=7304 km s^{-1}) and the H_I peak velocity at N1 (7295 km s^{-1}: the "systemic" velocity of the southern nucleus) (Baan, Hagiwara & Hofner 2007). These redshifted features arise from the southern nucleus but not from the northern nucleus. The tentative detection of the $V_{\rm LSR}$= 7158 km s^{-1} feature, which is blue-shifted by ∼150 km s^{-1} from the systemic, may constitute the first detection of counterpart to the redshifted features. Although NGC 6240 does not display the characteristic profile of maser emission from a rotating disk (with three distinct groups of maser features, one centered close to the systemic velocity and the others offset on either side by up to ∼1000 km s^{-1}), we note that if the red- and blue-shifted features are interpreted as originating in an edge-on disk (see, e.g., Miyoshi et al. 1995), the implied rotation speed ranges up to ∼ 300 km s^{-1}. Accordingly, the possibility of star-forming maser in NGC 6240 is less likely despite the low isotropic luminosity of the maser emission.

As already discussed in Hagiwara (2010), the non-detection of the maser toward the northern

TABLE 1
FLUX DENSITIES OF THE MASER AND DOUBLE NUCLEI IN NGC 6240

	Maser ($V_{\rm LSR}$, km s^{-1})		22 GHz continuum			
	7563.2	7609.3	Southern Nucleus		Northern Nucleus	
			F_P	F_T	F_P	F_T
(1)	(2)	(3)	(4)	(5)	(6)	(7)
2012 Jul 21	6.0±1.5	9.7±1.5	9.9±0.5	13.4±0.7	4.1±0.2	5.5±0.3
2012 Aug 28	<2.7[a]	5.1±0.9	7.7±0.4	11.9± 0.6	3.5±0.2	5.4±0.3

NOTE.—Col. (1): observing epoch; col. (2),(3): 22 GHz flux density (mJy) of each maser feature; col. (4)–(7): 22 GHz continuum peak flux (mJy beam^{-1}) and integrated flux density (mJy) of the southern nucleus (S) and northern nucleus (N)

[a]3 σ upper limit value

nucleus is not surprising even if both nuclei host nuclear masers, because of the very small possibility of the the disks around both nuclei in the galaxy being aligned edge-on to our line of sight.

4.2. M51

One of the most important results in our observations is that the 22 GHz nuclear continuum has been detected at the location of the radio nucleus of M51. This is the first detection of the nuclear continuum source at a frequency higher than 8 GHz, due to the improved VLA sensitivity. Moreover, we find that the location of the maser components coincides with that of the nuclear continuum. This indicates that the maser in M51 is a nuclear maser that is associated with AGN activity, although the maser is classified as a "low-luminosity maser" in Hagiwara (2007). By adopting an 8 GHz flux density of 0.21 mJy that was obtained by VLA A in Table 3, the spectral index, α (where $S_\nu \propto \nu^\alpha$, with S_ν being the flux density at frequency ν), is ≈ 0.10, very flat, assuming no flux variation. According to single-dish measurements in literature the total flux density is 142±15 mJy at 22.8 GHz (Klein et al. 1984). The total flux density of ∼0.4 mJy in our 22 GHz continuum map is clearly much lower, indicating that the source is heavily resolved on these angular scales and that we are seeing only the nucleus of the galaxy.

It is interesting to note that the jet axis that is associated with the nuclear continuum image is consistent with that previously reported at 5 and 8 GHz (Crane & van der Hulst 1992; Bradley et al. 2004), which confirms our detection and also shows that the velocity gradient detected in the earlier VLA data is seen along this jet axis. From our current data, there is no evidence that the M51 maser is a disk maser surrounding the nucleus of the galaxy at our resolution.

We failed to image the maser velocity gradient of Hagiwara (2010) in this data. The velocity gradient was later imaged by CO (1-0) and CO(2-1) emission at $\sim 0.4''$–$0.8''$ resolution in the nucleus of the galaxy, in good agreement with the velocity range and gradient direction of the maser (Matsushita et al. 2007), although these CO molecular gases are located $\approx 1''$ to the west of the maser or the radio nucleus. The velocity gradient in the previous data is real; however, we do not see the maser gradient in these observations, possibly due to the poorer (u, v) coverage resulting from the shorter observing time.

The weak off-nuclear blue-shifted emission

TABLE 2
A CUMULATIVE LIST OF H_2O MASER OBSERVATIONS IN NGC 6240

Epoch (yyyy.mm)	Telescope	Velocity range (km s^{-1} , LSR)	Maser velocities (km s^{-1}, LSR)	References
2000.03	Green Bank	7400−7700	7565.0±0.8	(1)
2001.01	Green Bank	6500−8100	7565.6±0.5	(1)
2001.05	Effelsberg	6850−7870	7565.0±1.1, 7609.0±1.1[a]	(2)
2001.06[b]	Nobeyama	6704−8858	7566.4±0.5	(3)
2001.12	Green Bank	6500−8100	7568.6±0.7	(1)
2002.04	Green Bank	6500−8100	7567.6, 7612.1±0.1	(1)
2002.06	VLA	7525−7665	7611.0±2.6[a]	(4)
2005.01	Nobeyama	6490−8610	7564.4±0.8	(5),(6)[c]
2007.01	Nobeyama	6490−8610	7442.0, 7561.5±0.8	(6)
2009.01	VLA	7370−7660	7491.1±0.2	(7)
2012.07	JVLA	6593−8321[d]	7563.2, 7609.3±0.8	This article
2012.08	JVLA	6593−8321[d]	7609.3±0.8	This article

·Notes.

[a]Uncertainties of velocity substituted by the channel spacings of Effelsberg or VLA

[b]Detected by averaging the spectra obtained in 2001 April and June

[c]By averaging the spectra obtained from 2003 to 2007, a narrow line maser feature centered at $V_{\rm LSR}$=7442.0±0.8 km s^{-1} was detected (K.Nakanishi et al. 2015, in preparation).

[d]Velocity range of −734 - 994 km s^{-1} to be added

References. — (1)Braatz et al. (2003), (2)Hagiwara et al. (2002), (3)Nakai et al. (2002), (4)Hagiwara et al. (2003), (5)Nakanishi et al. (2008), (6)K.Nakanishi et al. (2015, in preparation), (7)Hagiwara (2010)

TABLE 3
RADIO SPECTRA OF M51

Frequency: ν	5 GHz[a]	8 GHz[b]	22 GHz
S_ν[c] (mJy $beam^{-1}$)	0.89	0.21	0.23 ± 0.04
Spectral index: α_{8-22} (8–22 GHz)		0.1	

[a]Crane & van der Hulst (1992)

[b]Bradley et al. (2004)

[c]Peak flux densities measured by the VLA in the A-configuration

peaking at $V_{\rm LSR}$= 445 km s^{-1} that was detected in the earlier VLA run was not detected in our observation. The blue-shifted maser is not in the nucleus and if the excitation of the maser is due to star-forming activity, it would not be surprising that the maser is weak and variable (e.g., Baudry & Brouillet 1996).

5. SUMMARY

We report a correlated variation in the strength of the H_2O maser and the 22 GHz nuclear continuum emission in the southern nucleus of NGC 6240. The H_2O maser in the galaxy is additionally co-located with the nucleus and we conclude the maser is associated with AGN activity and not star formation. Thus, a scenario that argues for the possibility of a star-forming maser in the galaxy has been ruled out by our new observations. Several new minor H_2O maser features are detected at low significance toward the southern nucleus of NGC 6240 in the second epoch. More sensitive observations are required to confirm these features.

We also report the first detection of 22 GHz nuclear radio continuum emission in M51. The continuum emission is co-located with the maser position, which indicates that the maser arises from nuclear AGN activity and not from star-forming activity in the galaxy.

The extragalactic H_2O masers in both NGC 6240 and M51 show low isotropic luminosity on the order of 1 $L_\odot$, which is by two magnitudes lower than that of the water "megamaser," although it is most likely that the low-luminosity H_2O maser in the galaxies is a nuclear maser. The maser most likely amplifies the background radio continuum emission from the host galaxy nucleus. VLBI observations at milliarcsecond angular resolution would be able to reveal the distribution of the maser emission.

We thank the anonymous referee for useful comments on our manuscript. This research was supported by Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research(B) (Grant Number: 15H03644). This research has made extensive use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory (JPL), California Institute of Technology, under contract with NASA.

REFERENCES

Baudry, A. & Brouillet, N. 1996, A&A, 316, 188

Baan, W. A., Hagiwara, Y., & Hofner, P. 2007, ApJ, 661, 173

Beswick, R. J., Pedlar, A., Mundell, C. G. & Gallimore, J. F. 2001, MNRAS, 325, 151

Bradley, L. D., Kaiser, M. E., Baan, W. A. 2004, ApJ, 603, 463

Braatz, J. A., Wilson, A. S., Henkel, C., Gough, R., & Sinclair, M. 2003, ApJS, 146, 249

Carral, P., Turner, J. L., Ho, P. T. P., 1990, ApJ, 362, 434

Colbert, J. M. E., Wilson, A. S. & Bland-Hawthorn, J. 1994, ApJ, 436, 89

Crane, P. C., van der Hulst, J. M., 1992, AJ, 103, 1146

Depoy, D. L., Becklin, E. E., Wynn-Williams, C. G. 1986, ApJ, 307, 116

Dumas, G., Schinnerer, E., Tabatabaei, F. S., Beck, R., Velusamy, T., Murphy, E. 2011, AJ, 141, 41

Gallimore, J. F. & Beswick, R. J. 2004, AJ, 127, 239

Genzel, R., et al. 1998, ApJ, 498, 579

Greenhill, L. J., et al., 2009, ApJ, 707, 787

Hagiwara, Y., Henkel, C., Menten, K. M., Nakai, N. 2001, ApJ, 560, L37

Hagiwara, Y., Diamond, P. J., & Miyoshi, M. 2002, A&A, 383, 65

Hagiwara, Y., Diamond, P. J., & Miyoshi, M. 2003, A&A, 400, 457

Hagiwara, Y. 2007, AJ, 133, 1176

Hagiwara, Y. 2010, AJ, 140, 1905

Hagiwara, Y., Baan, W. A., & Klöckner, H.-R. 2011, AJ, 142, 17

Hagiwara, Y., Miyoshi, M., Doi, A., Horiuchi, S. 2013, ApJ, 768, L38

Heckman, T. M., 1980, A&A, 87, 152

Henkel, C., et al. 1984, A&A, 141, L1

Herrnstein, J. R., et al. 1999, Nature, 400, 539

Humphreys, E. M. L., Greenhill, L. J., Reid, M. J., Beuther, H., Moran, J. M., Gurwell, M., Wilner, D. J., Kondratko, P. T. 2005, ApJ, 634, L133

Ho, P. T. P., et al. 1987, ApJ, 320, 663

Klein, U., Wielebinski, R., & Beck, R. 1984, A&A, 135, 213

Komossa, S., Burwitz, V., Hasinger, G., Predehl, P., Kaastra, J.S. & Ikebe, Y. 2003, ApJ, 582, L15

Matsushita, S., Muller, S., and Lim, J. 2007, A&A, 468, L49

Miyoshi, M. Moran, J., Herrnstein, J., Greenhill, L., Nakai, N., Diamond, P., Inoue, M. 1995, Nature, 373, 127

Nakai, N., Sato, N., & Yamauchi, A. 2002, PASJ, 54, L27

Nakanishi, K., et al., 2008, in Mapping the Galaxy and Nearby Galaxies, Astrophysics and Space Science Proceedings, ed. K.Wada & F.Combes (Springer, New York), 364

Perley, R. A., Chandler, C. J., Butler, B. J., & Wrobel, J. M. 2011, ApJ, 739, L1

Reid, M. J., Honma, M. 2014, ARA&A, 52, 339

Risaliti, G., et al. 2006, ApJ, 637, L17

Sanders, D. B., Soifer, B. T., Elias, J. T., et al. 1988, ApJ, 325, 74

Scoville, N., & Young, J. S. 1983, ApJ, 265, 148

This 2-column preprint was prepared with the AAS LaTeX macros v5.2.



Fig. 1.— Comparison of the 22 GHz H_2O maser emission in the southern nucleus from 2-epoch VLA observations from the same velocity range. Left: the first epoch spectrum, observed in 2012 July. The maser features peaked at $V_{\rm LSR}$=7563.2 km s^{-1} and $V_{\rm LSR}$=7609.3 km s^{-1} (Gaussian-fitted) were seen. Right: the second epoch spectrum, obtained in 2012 August. The $V_{\rm LSR}$=7609.3 km s^{-1} was seen, while the $V_{\rm LSR}$=7563.2 km s^{-1} feature faded away in this epoch.



Fig. 2.— Tentative detection of the 22 GHz H_2O maser that consists of narrow features peaked (left) at $V_{\rm LSR}$=7158 km s^{-1}, (right) at $V_{\rm LSR}$=7372 km s^{-1}, $V_{\rm LSR}$=7396 km s^{-1}, and a broad feature ($V_{\rm LSR}\approx$ 7360 - 7400 km s^{-1}) toward the southern nucleus, all of which were obtained in the second epoch. No features are detected at the systemic velocity (7304 km s^{-1}). The peak velocities were Gaussian-fitted values. The detection needs to be confirmed.



Fig. 3.— 22 GHz continuum image of NGC 6240 (256 MHz bandwidth), obtained from the single-epoch VLA observation in 2012 July. The synthesized beam (FWHM) is plotted in the bottom left corner of the image. The contour levels are –4, 4, 5.6, 8, 11.3, 16, 22.6, 32, 45.2 of one σ value of 0.13 mJy beam^{-1}and the peak flux density is 7.75 mJy beam^{-1}.



Fig. 4.— Flux densities of the H_2O masers (filled and open circles) and 22 GHz continuum of the southern nucleus (open square), measured by the VLA in two epochs in 2012, are presented.



Fig. 5.— H_2O maser spectrum and continuum image of M51, observed in 2012 November, by the VLA in the A configuration are presented. Left: 22 GHz H_2O maser spectrum between $V_{\rm LSR}$= 460– 670 km s^{-1} toward the center of the galaxy. Right: 22 GHz continuum image of the galaxy obtained by using 256 MHz bandwidth. The contour levels are 3,4,5,6 of one σ value of 0.0365 mJy beam^{-1} and the peak flux density is 0.231 mJy beam^{-1}. The position of the detected H_2O maser ($V_{\rm LSR}$=549, 563, 565, 576km s^{-1} and some other minor features) is marked by a cross.